

NEUROBLASTOMA IN MUSCULOSKELETAL SYSTEM, TUMOR HYPOXIA, HIF-1 AND CHEMORESISTANCE

21th Prague-Lublin-Sydney-St. Petersburg Symposium

5th September 2019, Humpolec


Helena Doktorová

INTRODUCTION

1. Neuroblastoma (NB)
 - Symptoms
 - Staging
 - High-risk NB: indicators, therapy and treatment
2. Hypoxia in tumors
3. HIF-1 α
4. Our work: HIF-1 in chemoresistance
5. Inhibition options
6. Results
7. Other factors involved in hypoxia-induced chemoresistance
8. Conclusions


NEUROBLASTOMA

- Common and a frequently deadly cancer, almost exclusively seen in children
- The prevalence is about 1 case per 70 000 live births. This incidence is fairly uniform throughout the world
- It accounts for about 7–8 % of all cancers in children and the most common cancer in infants
- Nearly 90% of cases are diagnosed by age 5 X rare over the age of 10 years



Neuroblastoma among children in Southern and Eastern European cancer registries: Variations in incidence and temporal trends compared to US, Georgakis M. K. et al. , 2017

- Solid embryonal tumor
- Can arise anywhere throughout the sympathetic nervous system
- Most common primary site is adrenal gland found in the abdomen
- May also begin in nerve tissue in the neck, chest, or spinal cord.



- The clinical course of patients with neuroblastoma is highly heterogeneous

SYMPTOMS

- The “enigmatic tumor”
 - broad spectrum of clinical presentation, biological features and prognosis that vary greatly depending on size, location, and spread of the tumor
- Widespread bone and bone marrow disease causes bone pain -> limping or irritability
- Bone marrow replacement
- Anemia, bleeding, or infection
- Presenting symptoms of neuroblastoma depend on the location of the primary tumor, presence of metastatic lesions
- Weight loss and fever
- Metastases:
 - lymph nodes, bone marrow, bones, liver, skin
 - rarely to lungs or brain
 - multiple bone metastases in the vertebrae, metaphysis of long bones and skull

NB STAGING AND PROGNOSIS

- The **International Neuroblastoma Risk Group Staging System (INRGSS)** - results from imaging tests (such as CT or MRI and MIBG scan)
- The **International Neuroblastoma Staging System (INSS)** - results from the surgery

Stage 1:	Localized tumor* with complete gross excision, with or without microscopic residual disease; representative ipsilateral lymph nodes negative for tumor microscopically (nodes attached to and removed with the primary tumor may be positive).
Stage 2A:	Localized tumor with incomplete gross excision; representative ipsilateral nonadherent lymph nodes negative for tumor microscopically.
Stage 2B:	Localized tumor with or without complete gross excision, with ipsilateral nonadherent lymph nodes positive for tumor. Enlarged contralateral lymph nodes must be negative microscopically.
Stage 3:	Unresectable tumor infiltrating across the midline** with or without regional lymph node involvement; or localized unilateral tumor with contralateral regional lymph node involvement; or midline tumor with bilateral extension by infiltration (unresectable) or by lymph node involvement.
Stage 4:	Any primary tumor with dissemination to distant lymph nodes, bone, bone marrow, liver, skin and/or other organs (except as defined in Stage 4S).
Stage 4S:	Localized primary tumor (as defined for Stage 1, 2A, or 2B), with dissemination limited to skin, liver, and/or bone marrow*** (limited to infants less than one year of age).

The INRGSS has 4 stages:

L1	Localized tumor not involving vital structures as defined by the list of image-defined risk factors ^b and confined to one body compartment
L2	Locoregional tumor with presence of one or more image-defined risk factors ^b
M	Distant metastatic disease (except stage MS)
MS	Metastatic disease in children younger than 18 months with metastases confined to skin, liver, and/or bone marrow

← The INSS


HIGH-RISK TUMOR INDICATORS

Myc proteins:

- *MYCN* – amplification of *MYCN* oncogene is an established marker indicating aggressive tumor progression of NB
- c-Myc

DNA ploidy / DNA index:

- the higher the better prognosis in children under 2 years of age



Ref. Wang et al., *Diagnostic pathology*, 2013.: Representative FISH image of neuroblastic tumor cells displaying MYCN gene status. A: Amplification: The number of MYCN signals (green) is more than 10 copies of the CEP2 probe signals (red). C: No Alteration: Cells with MYCN signals (green) showing the same numbers of the CEP2 probe signals. (DAPI counterstain, original magnification $\times 1000$).

THERAPY OF HIGH-RISK NB

- Multi-agent chemotherapy
- Radiation therapy
- Stem-cell transplantation
- Surgery (if possible)

Problems in treatment:

- Late diagnosis
- Most NBs are radioresistant
- Chemoresistance



AFTER TREATMENT

- only 40–50 % survive 5 years after diagnosis
- NB can cause long-lasting side effects:
 - problems with bones and muscles – scoliosis, osteoporosis
 - chronic pain
 - problems with growth and development
 - metastases, bone marrow infiltration





FIG. 578. - Radiograms of the same case as Fig. 577. (a) Metastasis in the proximal humerus. (b) Calcification in retroperitoneal primary tumor.

HYPOXIA IN TUMORS

- The uncontrolled proliferation results in an imbalance between tumor cell oxygen consumption and tumor angiogenesis -> most parts of solid tumors are hypoxic
- Most tumor cells have adapted to these conditions
- Dramatic shift towards anaerobic metabolism -> increased expression of genes coding for glycolytic enzymes and glucose transporters
- Pro-angiogenic growth factors (VEGF, PDGF, angiopoietins)





➤ Tumor hypoxia is the main driving force behind vascularization and neoangiogenesis

➤ Metastases

Hypoxia inducible factor-1 α / HIF-1 α

- Hypoxia-induced **α subunit** creates the transcriptional complex with the constitutively expressed **β /ARNT subunit** -> HIF-1 transcription factor
- Normoxic degradation
- Hypoxic stabilization



HIF-1 regulates the expression of over 100 of genes that mediate the adaptive response of tumor cells to hypoxia

Hypoxia inducible factor-1 α / HIF-1 α

- Tab.1 Classification of the examples of HIF-1 regulated genes (Ref. Doktorova H. et al., Biomedical Papers, 2013)

BIOLOGICAL FUNCTIONS	GENE SYMBOL/ALIAS
Transcription factors	TWIST1, Snail, ZEB1, ZEB2, ID2, SMAD7, PPAR γ , GATA1
Histone modifiers	JMJD2B, JMJD2C, MLL1
Enzymes	MMP1, MMP3, LOX, ADAMTS1, ACE, ACE2, XPA, HK1, HK2
Receptors, receptor-associated kinase	CXCR4, CX3CR1, uPAR, PAI-1, 67-kDa laminin receptor, c-Met
Small GTPases, intracellular signalling molecules	VEGF , TGF- α , TGF- β 3, IGF2, Cdc42, Rac1, RhoE, IRS-2
Transporters	glut-1 , glut-3 , MDR1, VDAC1, transferrin, ceruloplasmin, IGF-BP1 - 3
Membrane proteins	ANGPTL4, L1CAM, α 5 integrin, CD151, CD24, CD147, Galectin-1 and 3, MUC1, Semaphorin 4D, Caveolin-1
Scaffold protein, cytoplasmic protein	HEF1, Lipirin- α 4
Matricellular proteins	CYR61, NOV


- There were described also several mechanisms in some types of tumors that regulate the stability and activity of HIF-1 in an oxygen-independent manner
 - > consequence of growth factor signalling
 - > in cells with defects in pVHL

Hif1a (hypoxia inducible factor 1, alpha subunit):

Gene expression/activity chart (mouse), physiological state

<http://biogps.org>

- Umbilical cells
- T- cells
- Follicular B-cells
- Mastocytes, Mast cells IgE
- Granulocytes
- 3T3-L1 preadipocytes
- Retina
- Bone marrow macrophage




Our work: HIF-1 and chemoresistance

- The overexpression of HIF-1 in cells results in higher resistance to the therapy and poor prognosis
- Targeting the HIF-1 pathway is as an important area for cancer therapy research
- We focus on *in vitro* study of high-risk neuroblastoma derived cell lines and their hypoxia-induced resistance to commonly used cytostatics, which is a major cause of anti-tumor failure in these children
- We have reported resistance of NB cell lines to cisplatine and ellipticine
- Other experiments: resistance to other commonly used cytostatics like doxorubicin, etoposide, melpahalan, 5-flouoruracil, gemcitabine and docetaxel

Inhibition options

- small molecule inhibitors (YC-1, LW6, Topotecan, Bisphenol A, Vorinostat etc. – different mechanisms of inhibition)
- gene knockdown, e.g. RNA interference (siRNA, shRNA)



Results

- HIF-1 inhibition is responsible for the growth suppression
(used cell lines: UKF-NB-4, Kelly, SK-N-AS)
- Lower hypoxia-induced resistance to cisplatin after HIF-1 inhibition
- Other factors have to be responsible for hypoxia-induced chemoresistance in NB cells




Fig. UKF-NB-4 cells were treated 48 h with 20 μ M cisplatin (CDDP) in normoxia or hypoxia (1 % O₂) and then analyzed by alamarBlue® - Cell Viability Assay. Cells were transfected with non-coding siRNA (NC siRNA) or HIF-1 siRNA, 24 h prior treatment. Similar results were observed in other studied NB cell lines. HIF-1 silencing has been verified by flow cytometry and Western blot analysis (data not shown).

Other factors involved in hypoxia-induced chemoresistance

- HIF-1 α independent
- factors that can somehow cooperate with HIF-1

Tab.2 Examples of other factors participating on chemoresistance due to the hypoxic conditions in cells

Anti-apoptotic factors (IAP3, Bcl-2 family)	HIF-1 independent
Cyclooxygenase-2 (COX-2), sphingosine kinase 2 (SphK2)	HIF-1 independent
Pim serine/threonine kinase-1 (Pim1)	HIF-1 independent
Nuclear factor kB (NF-kB)	can interact with HIF-1
Protein survival kinases PI3K and Akt	both dependent or independent on HIF-1
Signal transducer and activator of transcription 3 (STAT3)	both dependent or independent on HIF-1
Carbonic anhydrases (CA IX)	acts via HIF-1 pathway
V-ATPase	stabilises HIF-1 α in normoxia

CONCLUSIONS

- Cancer is still the leading cause of death. Prompt medical attention and aggressive therapy in high-risk NB treatment are important for the best prognosis.
- Continuous follow-up care is essential:
 - Side effects of radiation and chemotherapy, recurrence of the disease.
- New methods are continually being discovered to improve treatment and to decrease side effects.
- The reason for unsuccessful chemotherapy is the hypoxic environment:
 - Hypoxia-induced chemoresistance has been reported in a number of experiments
 - HIF-1 inhibition may have an impact on more efficient treatment of malignant tumors, it can suppress the growth of NB cells and their chemoresistance
- The detection of other factors can provide important prognostic features allowing subsequent individualization of anticancer therapy
- We still have more and better opportunities in research and so, there is hope...

Team and support: Hrabeta Jan, Groh Tomas, Mohamed Abdel Rahman,
Feriancikova Barbara, Cerna Tereza, Podhorska Natalia, Prof. Eckschlager Tomas

Department of Pediatric Hematology and Oncology, 2nd Medical School, Charles
University and University Hospital Motol, Prague, Czech Republic

Department of Biochemistry, Faculty of Science, Charles University, Prague, Czech
Republic



Thank you for your attention!